



Founded as a charity in 1999, the **Barrett's Oesophagus Campaign**, incorporating the McCord Fund, is the only national charity dedicated to the prevention of cancer from Barrett's Oesophagus and the support of people living with the condition.

We aim to:

- ⊙ provide a support and education service to people who suffer from Barrett's Oesophagus and their relatives and friends
- ⊙ support research to better understand the condition
- ⊙ raise awareness to encourage early detection

If you would like to find out more about the Campaign's work, or the services they offer, please contact:

The Barrett's Oesophagus Campaign
UCL, Division of Surgery and Interventional Science
Royal Free and University College Medical School
Rowland Hill Street
London NW3 2PF

Tel: 020 7472 6223
Fax: 020 7472 6224
Email: info@barrettscampaign.org.uk
www.barrettscampaign.org.uk

Registered Charity No. 1127495

working to prevent cancer of the gullet

design: www.worksong.co.uk



Treatments for
Barrett's
Oesophagus?

www.barrettscampaign.org.uk

This leaflet describes the various ways in which Barrett's Oesophagus is treated. There are two aims in treating Barrett's Oesophagus: to relieve the symptoms of acid reflux and to prevent it developing into cancer.

Treatment of acid reflux

People with Barrett's Oesophagus often have bad acid reflux but, curiously, may have very few symptoms. The treatment for reflux in people with Barrett's Oesophagus is the same as for people who do not have Barrett's. There are three approaches:

1. Things you can do for yourself

- avoid eating large meals and avoid eating within two hours of going to bed.
- The following foods tend to make reflux symptoms worse: chocolate, coffee, alcohol, fizzy drinks, spicy foods, citrus.

However, lifestyle changes will only help about one in five who have the condition.

2. Drugs

Antacids immediately neutralise the acid that has already been made. They may be either liquids or tablets and should be taken as

soon as you get symptoms.

Rennies and Tums, and most of the other medicines which you can buy over the counter, work in this way.

Alginates also contain antacids but, in addition, have a special ingredient which coats the lining of the stomach and oesophagus. This barrier prevents the acid from reaching the area where it would otherwise cause damage. Gaviscon and Gastrocote are examples of this class of medicine.

Acid suppression tablets work to stop acid being made before it can cause damage. There are two types: histamine receptor antagonists like ranitidine (zantac) and proton pump inhibitors (PPIs) such as omeprazole, lansoprazole, pantoprazole, rabeprazole and esomeprazole. These PPIs are far more effective at controlling acid reflux. Most patients with Barrett's Oesophagus will be taking one of these routinely.

3. Surgery

Fundoplication surgery is a treatment which aims to restore the normal valve at the lower end of the oesophagus which often does not work properly in people with Barrett's Oesophagus. This treatment is often carried out as a keyhole operation. You would only need to stay in hospital for one or two days, although it usually takes four weeks to recover completely from the operation.

Fundoplication surgery is successful in stopping acid reflux in seven or eight of every ten people treated. It does have risks, so before agreeing to have surgery it is important to discuss these with the surgeon. Things which can trouble people after surgery include bloating of the abdomen, difficulty in swallowing and, rarely, severe diarrhoea.

When to use which treatment

Everyone with acid reflux should try to make lifestyle changes. Often, simple changes to when and what they eat will make people feel very much better.

Many people will still need to take drugs. People with frequent or

severe symptoms should consider taking acid suppression tablets to try and prevent complications of acid reflux such as scarring of the oesophagus.

Surgery is usually reserved for people who do not respond well to lifestyle changes and drug therapy.

Treatments to prevent cancer

Since the vast majority of patients with Barrett's Oesophagus do not get cancer, the usual practice in the United Kingdom is not to attempt to remove the Barrett's cells. Treatment is usually only offered if the cells look as though they are starting to change and the risk of getting cancer starts to rise.

Acid suppression and fundoplication surgery

Although in theory, exposure to acid and bile may make cells more likely to turn cancerous, there is no clear evidence that aggressive suppression of acid reflux does actually reduce the risk of cancer. Decisions about these treatments should generally be made on the basis of symptoms.

Dysplasia

This word is derived from the Greek meaning roughly “bad formation”. It is used to describe an abnormality within a tissue where the tissue changes and may in some cases progress to cancer. Dysplasia is the earliest form of pre-cancerous change that can be recognized and maybe low grade or high grade, the latter representing a more advanced progression towards cancer.

Treatment for people with low grade dysplasia

The risk of people with low grade dysplasia getting cancer is only slightly higher than in people without dysplasia. For this reason most doctors do not recommend treatment for this condition.

Most doctors would recommend that you have check-ups with an endoscopy every six months so that they can monitor any changes in the Barrett’s cells. This can be done at your local hospital. This is the current guideline of the British Society of Gastroenterology.

Treatment for people with high grade dysplasia

The risk of developing cancer is much higher for people with high

grade dysplasia, at about 10 per cent per year, with half of all patients developing cancer in around five years. This means that half of these people will NOT develop cancer in five years. This needs to influence your decision about whether or not to have treatment.

All people with high grade dysplasia should be referred to a specialist centre where their case is discussed by a multi-disciplinary team of doctors and where the doctors perform minimally invasive but complex treatments frequently. All these treatments aim to remove the dysplasia. In addition, they may aim to replace the red, Barrett’s Oesophagus, with normal pink (squamous) lining.

Endoscopic Mucosal Resection

Some patients with high grade dysplasia have a visible nodule in their oesophagus. It is relatively straightforward to remove the nodule during endoscopy. The procedure takes around 30-45 minutes and you can usually go home the same day. If you have this procedure you will be given a sedative to make you slightly

sleepy. Most people can eat and drink normally afterwards. In about one in ten people there may be minor bleeding, and more serious bleeding in one in 50 people.

This procedure can be repeated a number of times if there are several nodules, but it cannot remove large sections of affected oesophagus without causing scarring and difficulty in swallowing.

Endoscopic mucosal resection is a particularly useful technique if the diagnosis is not clear because the removed nodule can be sent to the laboratory to be checked by the pathologist. In this situation, it serves as both a diagnostic test and a treatment. This treatment does not aim to remove the Barrett’s Oesophagus cells completely.

Photodynamic therapy (PDT)

This treatment is particularly useful when there is high grade dysplasia but no nodules. Here the changes are often widespread and are difficult to see by endoscopy. PDT can be used to treat a large area.

This treatment does not aim to completely remove the Barrett’s Oesophagus although it sometimes does so.

If you have PDT, you will be given a drug which sensitises you to light. You then have an endoscopy, during which light is shone at the area which needs treatment.

The combination of the drug and the light kills the targeted cells. This treatment takes about 45 minutes. The treatment can be repeated two or three times at an interval of three months if necessary and has been shown to reduce the likelihood of cancer developing by 50 per cent over at least five years.

The treatment, using a drug called Photobarr, is licensed and has been approved by NICE (the UK government committee regulating the use of new treatments) for treating high grade dysplasia in Barrett’s Oesophagus.

For two to three weeks after your treatment you will probably have some discomfort when you swallow but this then resolves. In addition, the drug used, Photobarr, will cause your skin to be sensitive to light for up to three months.

Take care to avoid bright sunlight during this time. A number of other drugs are available for PDT but these are only used in studies and are not widely available.

Around one in four people will develop scars in the oesophagus after treatment which make swallowing difficult. This can be treated by stretching the oesophagus during another endoscopy. Some people need to have this done several times to return their swallowing to normal.

HALO radiofrequency ablation

This is a new treatment which does not have the risks of skin sensitivity to light. It uses radio waves to destroy the dysplasia.

You will be given a sedative to make you sleepy. A tiny probe will be used during an endoscopy to deliver the radio waves to the affected parts. The procedure takes about 45 minutes. Some people return to normal immediately after treatment, but many feel nauseous and have chest pain, particularly when they eat, for up to three weeks after treatment. A very few people (around 1 in 100) suffer scarring of the oesophagus. This

treatment is usually repeated three or four times at intervals of two to three months until not only the dysplasia, but also the entire Barrett's Oesophagus has been removed.

At present, the outcomes of this treatment look very promising, but as yet there is not the information about how long the benefits last.

Research is going on all the time into new ways to treat Barrett's Oesophagus. New studies are being published regularly. Please speak to your specialist about the current state of knowledge regarding the treatments available.

The facts:

In the UK:

- ⊙ at least 375,000 people are estimated to have Barrett's Oesophagus
- ⊙ 37,500 of these can be expected to develop oesophageal cancer
- ⊙ up to 30,000 can be expected to die within five years